Hand Foot and Mouth Disease (HFMD)

West Moreton Public Health Unit has been advised of a case of hand foot and mouth disease at your school/child care centre.



- a mild illness
- easily spread in schools and child care settings
 not related to foot and mouth disease in animals
 can (rarely) lead to a more serious illness



- painful tiny blisters or ulcers in or on the:

 - mouth,
 mouth,
 tongue,
 palms of hands, fingers,
 soles of feet,
 legs,
 nappy area.
- fever

- feeling unwell
 painful eating and drinking
 blisters can be sore, itchy
 and last up to 10 days
- can be peeling of the hands and feet up to 3 weeks after infection







- Contact with fluid from
- the blisters
 contact with saliva, drool,
 nasal mucous, faeces (poo)
 contaminated surfaces





No specific treatment is needed for HFMD.

- Manage fever and discomfort with paracetamol or lbuprofen.
 Make sure children get enough fluids cold and soft food might be easier to swallow.



Children with symptoms of hand foot and mouth disease should stay at home until feeling well and all the blisters have dried



REPORT

Let the child care centre or school know your child may be infectious with hand foot and mouth disease



WASH HANDS well and often using soap especially, • before eating or handling food • after using the toilet • changing nappies, soiled clothes • contact with infected skin





COVER COUGHS and SNEEZES,

teach children how to cough/sneeze into their elbow or use a tissue and wash hands afterwards



- CLEAN and DISINFECT,
 frequently touched surfaces
 soiled items including toys
 avoid sharing cups, eating utensils and personal hygiene items (towels, toothbrushes, wash cloths) or clothing (shoes and socks)



- you are worried your child is not eating or drinking has a severe headache
- · has persistent fevers

For more information see the QH fact sheet at:

West Moreton Public Health Unit

© 07 3271 8744

13HEALTH (13 43 25 84)





